

## Follicle Wave and Natural Estrus Synchronization Superovulation in Holstein Cows

Manuel E. Soria Parra; Carlos A. Soria Parra; Daniel Argudo Garzón; Guillermo Serpa García; Silvana Méndez Álvarez; Carlos Torres Inga; Guillermo E. Guevara Viera, PhD.

Faculty of Agricultural Sciences, University of Cuenca, Yanuncay Campus, Ecuador

manuel.soria@ucuenca.edu.ec

### ABSTRACT

The effects of two superovulation protocols were compared (synchronization of the follicle wave SFW and natural estrus NE) in embryos collected from Holstein cows. Twenty cows were chosen as donors, body condition of 2.75 - 3.5; between 40 and 60 months, with normal cycles and no breeding problems. The cows were randomly assigned to SFW treatment (n=10), and the NE treatment (n=10). The SFW group was synchronized with intravaginal progesterone plus estradiol benzoate, on day 0, and increasing doses of FSH for 4 days, twice a day, from the 4<sup>th</sup> on. The implant was removed on the 6<sup>th</sup> day, and prostaglandin was applied. On the 8<sup>th</sup> day, insemination was made at 6 am and 6 pm. The NE group received increasing doses of FSH twice a day, during 4 days, from the 10<sup>th</sup> day on. On the 12<sup>th</sup> day prostaglandin was administered, and insemination took place on the 14<sup>th</sup> day, at 6 am and 6 pm. The embryos were recovered from the two groups without surgery, 7 days after the first artificial insemination. The values of embryos for transference were  $5.7 + 0.76$  and  $2.8 + 0.31$  ( $P < 0.05$ ) for SFW and NE, respectively.

**Key words:** bovine, progesterone, estrus, prostaglandin

### INTRODUCTION

Understanding follicle dynamics and its self-regulated processes, is the bases of the new alternatives to control estrus cycle in cows. The generation of new knowledge about follicle dynamics in bovine ovaries helps reformulate the traditional methods of estrus control, generating new proposals of estrus and ovulation induction and synchronization.

Superovulation programs (SVP) have been developed in donor cows: based on natural estrus (NE), or by follicle wave synchronization (SFW) (String fellow and Seidel, 2000; Baruselli *et al.*, 2006; Bó, Tríbulo and Mapletoft, 2011).

Superovulation includes dealing with endocrine processes during the estrous cycle, until ovulation of follicles that grew as a response to hormones occurs. It is triggered by follicle growth waves to produce ovocytes which can be fertilized, and seven days later the embryos can be recovered and classified according to their stage or quality, as established by the International Embryo Transference Society (IETS) (Bochesta and Nasser; 2005; Belacuba, 2007; Amaya, 2010; Luca and Castrillón, 2011).

The purpose of this paper was to compare the effects of two superovulation protocols (synchronization of the follicular wave, SFW, and natural estrus, NE) in the quality of embryos collected from Holstein cows.

### MATERIALS AND METHODS

Research took place in the province of Azuay, Cuenca canton, parishes of Tarqui, Cumbe and Victoria del Portete, located on 20; 18 and 8 km from the city of Cuenca, respectively.

Twenty Holstein cows were chosen to make two 10-animal groups, based on the following parameters: body condition (2.75 - 3.5); 40 - 60 months old; normal estrous cycles, minimum of 60 open days, and no breeding problems, according to previous gynecological and ecographic assessments.

The following protocols were applied:

SFW group (n=10):

- Day 0: intravaginal implant (IV) of 1.38 g of progesterone (P4; CIDR®, Zoetis, Mexico), 2 mg of estradiol benzoate (EB; Gonadiol®, Syntex, Argentina) IM; and 100 mg of P4 (Gestavec® Vecol SA, Colombia) IM.
- Day 4: 40 mg of Folltropin® (Bioniche, Canada) IM twice a day (6 am and 6 pm).
- Day 5: 30 mg of FSH-p/IM twice a day (6 am and 6 pm).
- Day 6: 20 mg of FSH-p and 0,5 mg of D-chloprostenol (Estrumate® Intervet Schering-Plough, NJ, USA) IM twice a

day(6 am and 6 pm); ó IV was removed at 6 pm.

- Day 7: 30 mg of FSH-p/IM twice a day(6 am and 6 pm).
- Day 8: artificial insemination(AI) at 6 am and 6 pm, Gonadorelin (GnRH; Fertagyl®, Intervet Schering-Plough, NJ, USA) 500 ug/IM at 6 am.

NE group (n=10):

- Day 0: natural estrus.
- Day 10: observation of luteal body and absence of the dominant follicle, through transrectal echographic scanning and 40 mg of FSH-p/IM twice a day (6 am and 6 pm).
- Day 11: 30 mg of FSH-p/IM twice a day(6 am and 6 pm).
- Day 12: 20 mg of FSH-p and 150 ug of D-cloprostenol/IM twice a day(6 am and 6 pm).
- Day 13: 30 mg of FSH-p/IM twice a day(6 am and 6 pm).
- Day 14: IA at 6 am and 6 pm, GnRH 500 ug/IM at 6 am.

Seven days after the first AI in the two groups, the embryos were collected without surgery, using saline buffer solution (PBS).

The embryos were spotted and isolated in a Holding maintenance medium, using a diascopic light stereoscope (Vigro™, Bioniche, Pullman, WA, USA). Quality assessment was made according to the embryo classification parameters (IETS, 2000). The variables studied were total recovered structures (ovocytes and embryos), transferable embryos (quality embryos 1 and 2), and non-transferable embryos (quality embryos 3, degenerated embryos and non-fertilized ovocytes). The basic statigraphs were estimated, and the Student test was made to independent samples;  $P < 0.05$  was considered statistically significant.

## RESULTS AND DISCUSSION

The current protocols include estrogens (E2) and P4, which control the appearance of the follicle wave (Bó, Tribulo and Mapletoft, 2011), and the moment of ovulation (Baruselli *et al.*, 2006) have made possible the application of superovulation treatments (SVT) in donor cow groups, at any stage of estrus. Comparison of the SVT (which included NE), and SFW in terms of structures recovered, resulted in  $6.5 + 2.02$  and  $8.7 + 1.86$ , re-

spectively, with  $P > 0.05$  (see table). It corroborated the study made by Bó *et al.* (1996), where no significant differences were found concerning the amount of ovocytes and embryos recovered. It compared a superovulation treatment with progesterone + beta estradiol to synchronize the wave, to the control group that took estrus as reference. Similar results were found by Goulding *et al.* (1990), with  $8.5 + 0.9$  ovocytes and embryos in SVT, and initiated treatment with FSH on the 10<sup>th</sup> day of estrus.

Several studies (Bo *et al.*, 1995; Mapletoft, Bennett Steward and Adams, 2002) coincided in that starting a superovulation treatment requires the absence of a dominant follicle. It can be achieved with E2 + P4 to synchronize follicle wave, or initiate treatment with FSH-P between days 10 and 12 during the cycle. In both cases, emergence follicles with similar superovulation response were achieved; additionally, a similar amount of ovocytes and recovered embryos were generated, using the proper gonadotrophin dose were achieved.

Concerning embryo quality, a significant difference was observed between the transferable structures achieved by SFW, when the SOV treatment was made, taken NE  $5.7 + 0.76$  and  $2.8 + 0.31$  as reference ( $P < 0.05$ ), respectively (see table). These results match the parameters of the International Embryo Transfer Society (IETS, 2000), in relation to the fact that the structures recovered must be 40 - 50% transferable. Exogenous P4 applied to the SFW treatment could guarantee the generation of transferable structures. P4 is known to play a role in ovocyte quality. Salhab *et al.* (2011) claimed that P4 has an important effect on ovocyte quality because the cummulus cell apoptosis frequency decreases during maturation (Aller *et al.*, 2015).

Rivera *et al.* (2011) described the existence of a positive relationship between high blood P4 levels during follicle growth and embryo quality, in dairy cows. However, other authors, like Fair and Lonergan (2012), indicated that such effect has not been confirmed, but it might be related to a decline in ovocyte exposure to LH, thus preventing premature maturation (Cerr *et al.*, 2011). Synchronization of the follicle wave dependos on GnRH to facilitate development of a wave of follicles by the action of FSH and P4, which allow for maturation of ovocytes. Also, an increase in LH pulse frequency triggers ovulation.

Regarding non-transferable structures, the results were  $3.0 + 0.86$  and  $3.67 + 1.45$  for SFW and NE, respectively (see table). Jiménez (2009) said that there are external factors that affect superovulation response: species, breed, age, physiological state, fertility, characteristics of follicle waves, individuality, and others. It would determine that regardless of the superovulation treatment applied, there will always be non-transferable structures.

## CONCLUSIONS

The superovulation treatment based on SFW produces a greater amount of transferable embryos in comparison to the treatment based on NE. The SFW protocol has the advantage of better follicle wave due to ablation of the dominant follicle. It also permits synchronization of follicle wave. Besides, exogenous P4 contributed to more high quality oocytes. The addition of GnRH during insemination provided synchronization at ovulation, producing higher quality and quantity of transferable embryos.

## REFERENCES

- ALLER, J. F.; ABALOS, M. C.; ACUÑA, F. A.; VIRGILI, R.; REQUENA, F. y CANCINO, A. K. (2015). Embryo Yield In Llamas Synchronized With Two Different Intravaginal Progesterone-Releasing Devices and Superovulated With eCG. *Spanish Journal of Agricultural Research*, 13 (3), 3-8.
- AMAYA, A. D. (2010). *Efeto del diametro del foliculo ovulatorio, tamaño del cuerpo luteo y perfiles de progesterona sobre la tasa de preñez en la hembra receptora de embriones bovinos*, 1-190. Tesis en opción al título de Máster en Salud Animal, Bogotá, Colombia.
- BARUSELLI, P. S.; DE SÁ FILHO, M. F.; MARTINS, C. M.; NASSER, L. F.; NOGUEIRA, M. F. G.; BARROS, C. M. y BÓ, G. A. (2006). Superovulation and Embryo Transfer in *Bos indicus* Cattle. *Theriogenology*, 65 (1), 77-88. Retrieved on November 24, 2015, from <http://doi.org/10.1016/j.theriogenology.2005.10.006>.
- BELACUBA, F. (2007). *Factores que afectan la superovulación en bovino*. Retrieved on November 24, 2015, from [http://www.produccion-animal.com.ar/informacion\\_tecnica/transplante\\_embriionario/17-superovulacion.pdf](http://www.produccion-animal.com.ar/informacion_tecnica/transplante_embriionario/17-superovulacion.pdf)
- BO, G. A.; ADAMS, G. P.; CACCIA, M.; MARTÍNEZ, M.; PIERSON, R. A. y MAPLETOFT, R. J. (1995). Ovarian Follicular Wave Emergence After Treatment With Progestogen And Estradiol In Cattle. *Animal Reproduction Science*, 39 (3), 193-204.
- BO, G. A.; ADAMS, G. P.; PIERSON, R. A. y MAPLETOFT, R. J. (1996). Effect of Progestogen Plus Estradiol-17 $\beta$  Treatment on Superovulatory Response in Beef Cattle. *Theriogenology*, 45 (5), 897-910.
- BÓ, G. A.; TRÍBULO, A. y MAPLETOFT, R. J. (2011). Nuevos protocolos de superovulación para programas de transferencia de embriones en bovinos. *SPERMOVA*, 1(1), 26-33.
- BÓ, G., y NASSER, L. (2005). *Efficiency of Programs that Control Follicular Development and Ovulation for the Donor Superovulation without Estrus Detection*. Minneapolis, EE.UU.: Proc Joint Mtg Am Embryo Trans Assoc & Can Embryo Trans Assoc.
- CERRI, R.; CHEBEL, R.; RIVERA, F.; NARCISO, C.; OLIVEIRA, R.; THATCHER, W. y SANTOS, J. (2011). Concentration of Progesterone during the Development of the Ovulatory Follicle: I. Ovarian and Embryonic Responses. *Journal of Dairy Science*, 94 (7), 3342-3351.
- FAIR, T. y LONERGAN, P. (2012). The Role of Progesterone in Oocyte Acquisition of Developmental Competence. *Reproduction in Domestic Animals*, 47 (4), 142-147.
- GOULDING, D.; WILLIAMS, D. H.; DUFFY, P.; BOLAND, M. P. y ROCHE, J. F. (1990). Superovulation in Heifers given FSH Initiated Either at Day 2 or Day 10 of the Estrous Cycle. *Theriogenology*, 34 (4), 767-778.
- JIMÉNEZ, C. (2009). Superovulación: estrategias, factores asociados y predicción de la respuesta superovulatoria en bovinos superovulation: strategies, associated factors, and prediction of the superovulatory response in cows. *Revista de la Facultad de Medicina Veterinaria y de Zootecnia (Universidad Nacional de Colombia)*, 56 (3), 195-214. Retrieved on November 24, 2015, from <http://www.revistas.unal.edu.co/index.php/remvez/article/view/13769>.
- LUCA y CASTRILLÓN, M. (2011). Respuesta a la superovulación y calidad de los embriones en bovinos lecheros de elevado mérito genético, uso de diferentes protocolos. *engormix*. Retrieved on November 24, 2015, from <http://www.engormix.com/MA-ganaderia-carne/genetica/foros/respuesta-superovulacion-calidad-embriones-t22419/103-p0.htm>.
- MAPLETOFT, R. J.; BENNETT STEWARD, K. y ADAMS, G. P. (2002). Recent Advances in the Superovulation in Cattle. *Reproduction Nutrition Development*, 42 (6), 601-611.
- RIVERA, F. A.; MENDONCA, L. G. D.; LÓPEZ, G.; SANTOS, J. E. P.; PÉREZ, R. V.; AMSTALDEN, M., et al. (2011). Reduced Progesterone Concentration During Growth of the First Follicular Wave Affects

Embryo Quality but Has No Effect on Embryo Survival Post Transfer In Lactating Dairy Cows. *Reproduction*, 141 (3), 333-342.

Salhab, M.; Tosca, L.; Cabau, C.; Papillier, P.; Perreau, C.; Dupont, J.; *et al.* (2011). Kinetics of Gene Expression and Signaling in Bovine Cumulus Cells Throughout IVM in Different Mediums in Relation

to Oocyte Developmental Competence, Cumulus Apoptosis and Progesterone Secretion. *Theriogenology*, 75 (1), 90-104.

STRINGFELLOW, D. A. y SEIDEL, S. M. (2000). *Manual de Transferencia de Embriones* (3ra ed.). Sociedad Internacional de Transferencia de Embriones.

Received: 6-22-2016

Accepted: 7-1-2016

**Table. Results from embryos in treatments with SFW and NE**

<i>Structures</i>	<i>Estructuras</i>	SFW group (n=10)	NE group (n=10)	P value
Recovered (ovocytes and embryos)		8.7 ± 1.86	6.5 ± 2.02	NS
Transferable embryos (Quality 1 and 2)		5.7 ± 0.76	2.8 ± 0.31	*P < 0.05
Non transferable (Quality 3, degenerated and non-fertilized oocytes)		3.0 ± 0.86	3.67 ± 1.45	NS

NS indicate insignificant differences  
NE: natural estrus

SFW: synchronization of follicle wave