

## Composition, Attributes and Benefits of Goat Milk: Literature Review

Adela Bidot-Fernández

Center for Animal Breeding Research in Tropical Livestock Raising (CIMAGT)

abidot@infomed.sld.cu

---

### ABSTRACT

Goat's milk and cow's milk have similar compositions. The former is healthy and nutritious, and it is a valid alternative to human's milk because their nutritional values are also very similar. Many people who are allergic to cow's milk can drink goat's milk safely, as it contains a different protein. Goat's milk is gradually becoming more popular in world markets, beyond the countries where it is already one of the major components in the diet of millions of people today. To produce good quality milk some basic efficiency principles of livestock farming must be taken into consideration, like good quality animals, selection of dairy genotypes, adequate food supply, good management and health. The first two have direct effects on the nutritional quality or composition of milk; the other two have to do with hygiene. This paper reviews the main aspects that influence goat's milk composition, biological value as substitute for human's milk, main components, and qualities and benefits to human health.

**Keywords:** *goat's milk, milk composition, quality of milk*

### INTRODUCTION

Goat was the first animal domesticated by man that could produce food for humans, about 10 000 years ago. Since then, goats were present along history, according to several historical, mythological and Biblical accounts that mention the caprines. However, few times had they been recognized (Doria, 1997, Bidot and Muñoz, 2016).

Goat milk has been used for human nutrition since ancient times. Very old records in the Bible or Egyptian murals referred to the consumption of goat's milk. Its history is linked to the history of man, who has always used the goats', meat, fur, leather, manure and labor. These products are important indicators of the species' capacity to adapt to multiple climates and systems (Cofré, 2001).

Most part of the caprine population, approximately 444 000 000 (95%), lives in developing countries, and they are used as dual-purpose (meat-milk) animals. However, the industrial countries only have 5% of goats (30 000 000), especially for milk production (27% of total world milk production) (FAO, 1982).

According to FAO (1987), goats are one of the few species that can survive and even produce food in adverse conditions, with harsh climates and scarce natural resources.

The goat's milk demand estimated by FAO by 2000 was 177.6 million tons, mostly from developing tropical countries (Knights and García, 1997). By 2010, FAO recorded 909 million of heads, whereas the countries with the largest numbers were India (154 million), China (150

million), Pakistan (59 million), and Sudan (43 million).

In the MERCOSUR countries, the caprine production combined accounts for 1.8% of the world's total: Argentina (4.2 million), Brazil (9.3 million), Paraguay (135 thousand), Chile (750 thousand), and Uruguay (16.7 thousand) (Bedotti, 2008; FAO, 2010).

Despite its important contribution to human nutritional support and other areas (clothing, labor, fertilization), goats have been reviled as an enemy of ecosystems and large contributors to the loss of cropland. Hence, they have been isolated to deserted and irregular places. Goats are also considered responsible for transmitting diseases to humans (brucellosis, Malta fever), and for spreading them worldwide through the large commercial and agricultural routes. Based on such pejorative considerations and other causes (geographic, social, economic, situational) caprine raising has undergone unequal evolution in different parts of the world (Vacas, 2003).

López *et al.* (2011) noted that the marginality of farm productions, especially caprines, is characterized by the absence of practices that provided added value to productions. The family rationale to increase income is to have large numbers of animals, instead of managing less and get higher productivity from them. Either will require proper technology according to the ecological and socio-economic circumstances, as well as the markets to sell.

Goat milk has been a key component in the Mediterranean diet originally, particularly for-

making cheese, as noted by the literature classics Caton, Virgilio, Columela, Plinius, and Atheneus. They not only showed how to make cheese, but also the kinds available ("oxigala", "moretum"), or even some recipes, like pie ("sabilium"), made of cheese, honey, flour and eggs, powdered with poppy seeds and baked in the oven (Otogalli and Testolin, 1991; Capdevila and Martí-Henneber, 1996). In ancient times, fermented milk was mentioned in the Bible's Deuteronomy, as "one of the foods given by the Lord to His people". Since then, goats have played a key role in quality foods for humans, especially in poverty-stricken areas of the world, where these foods continue to be the main source of proteins for the people (Bidot, 2006b). A tenth of all the milk consumed in the world is produced by goats; for some countries, it is the only source of milk (Arbiza, 1987).

Goat's milk demand has increased, mainly due to demographic boom, and the particular interest in goat's milk by-products in developed countries (yogurt, and cheese), because they can be consumed by groups of people who are lactose intolerant. Goat milk composition is associated to certain nutritional benefits to children, and the development of functional foods and by-products with particular sensorial features required by consumers. This food and its derivatives are another choice to invigorate regional economies (Arbiza, 1996; Haenlein, 2004; Vega and León *et al.*, 2010).

## DEVELOPMENT

### *Composition of goat milk (Table 1)*

From a technological perspective, milk composition determines its nutritional quality, properties and value as raw material to make foods. Goat's milk has the highest nutritional and therapeutical values, only second to human's milk, with a high nutritional quality and agreeable taste. The therapeutical properties of goat's milk were acknowledged since the beginning of civilization, proving its power against gastrointestinal ailments (Flores Cordova *et al.*, 2009).

Milk is the white-beige, fluid product secreted by the female's udder, with a particular odor and taste. It is rich in nutrients and easily contaminated if not properly collected. In general terms, goat milk is matt-white and slightly viscous, and its composition and physico-chemical features are very sensitive. Some of the factors that contribute

to these variations are, the breed, nutrition, season, environmental conditions, area, lactation conditions and udder health (Chilliard *et al.*, 2003; Park, 2007a; Park *et al.*, 2007).

The goat's milk is whiter than the cow's milk because it has no carotene that provide a yellow tone. Carotens are red, orange or yellow non-saturated hydrocarbons found in tomatoes, carrots, egg yols, and other plants. In animals, it is transformed in vitamin A. Goat's milk has a strong odor caused by the absorbtion of aromatic compounds during husbandry (usually inadequate substances), with the presence of males in the milking areas, poor housing hygiene, delayed skimming and after-milk cooling. The said odor and taste can be removed by vaccuum deodorization, which is very simple (Borras, 1968). Cow's milk is slightly acidic, whereas goat milk is almost alkaline (pH 6.7), due to higher protein contents and different phophate combinations (Saini and Gili, 1991), so people with digestive problems are encouraged to consume this kind of milk (Jandall, 1996). Goat's milk is a balanced combination of proteins, fats, carbohydrates, salts and other components. Milk composition determines its nutritional quality and its value as raw material to manufacture foods for humans. Its qualitative composition is stable, but it varies concerning the animal breed, the lactation time, number of deliveries, season, and local climate. Other authors describe milk as a white and opaque liquid with a complex composition, slightly sweet and an almost neutral pH. It is a suspension of protein elements within serum made by lactose and mineral salts, mainly (Alais, 1988; Ortega *et al.*, 2011).

According to Vargas *et al.* (2007), milk is a white, opaque liquid, twice as dense as water, slightly sweet and with little odor. It is a very complex chemical and physical system, and it can be considered an emulsion of fatty matter in an aqueous solution that contains various dilluted or colloidal elements.

According to the definition of milk, adopted by the I International Conference for Food Fraud Repression, in Geneva, 1908, it is "the product...colostrum..." It basically coincides with the definition established in the Spanish Nutritional Code (Real Decree 2484/1967, Spetember 21), Chapter V, which noted that milk is the whole product without alterations or forgery, without colostrum from regular, clean, complete, and unin-

errupted milking of healthy well-fed mammal females." (Vargas *et al.*, 2007).

A WHO report, Series of Technical Reports No. 124, Geneva, 76, 5, 1957, established that milk and dairy products which are produced without proper hygienic conditions may be the cause of diseases in the humans that consume it. Therefore, milk hygiene comprises dairy animal hygiene, application of adequate hygienic methods for production, handling and manufacturing of milk and dairy products, pasteurization, or other forms of heat treatments to destroy pathogenic germs, and the protection of the product against further contamination. The Committee resolved that its attention had to be focused on hygienic issues related to cow's milk (including buffalos), then goat's milk. Additionally, issues related to dairy products, like butter, cheese and ice cream would be considered by another Committee of Experts, called upon by WHO and FAO (Meneses, 2007).

Today, milk is gathering greater importance, especially in terms of protein percents. Milk rich in total solids can produce higher yields in dairies. The caprine dairy industry must have information on the milk collected and sent by suppliers over a year, and measure the physical and chemical parameters useful to accept or refuse the main product and pay the manufactures (Cruz *et al.*, 2012).

The composition of milk is 77-80% water, and 20-23% total solids. They are usually composed of 3-3.5% fat, 3-3.5% protein, and 4-6% of carbohydrates, like lactose and minerals (calcium) (Salvador *et al.*, 2006).

Nutritional facts of goat's milk (Pérez, 2001). In 100 grams of goat's milk there are,

Calories: 70 kilocalories

Carbon hydrates: 4.5 g

Proteins: 3.3 g

Fats: 4 g

Cholesterol: 11.0 mg

Glycemic index: 24

Vitamins: A, D and C; lower in vitamins B1, B2, B3, B5 and B12.

Minerals: calcium, phosphorous, potassium, magnesium, iron, zinc, selenium, manganese and copper.

Other authors compare cow's and human's milk to goat's, though there may be differences in terms of breeds and nutrition.

The goat's milk protein often has a range of essential and total aminoacids of 0.46, and of essen-

tials against nonessentials, of 0.87 (Singh and Singh, 1985). The size of casein mycells is lower in goat's milk (50 nm), compared to cow's milk (57 nm) Alais, 1988). The caseins in goat milk have larger glycine contents and less arginine and sulphur aminoacids, especially methionine (Capra, 2004).

The mineral contents of goat's milk is greater than human's milk. Goat's milk contains about 134 mg of Ca and 121 mg of P, per 100 of milk. It can even have up to 13% more calcium than bovine's milk, but it is not as good a source of iron, cobalt and magnesium. Table 4 shows the values reported for the amounts of minerals found in goat's and cow's milk (Park, 2006).

In short, milk must be of good quality for consumption as milk, or as by-products, which means that besides having high nutrient contents, it must have special features that guarantee fresh, nutritional and healthy products on the market (Rodríguez and Valencia, 2006).

#### *Properties*

Goat's milk is close to perfection, with an amazing structure, very similar to human's milk. The differences in many cases mean a broad number of nutritional advantages for goat milk over many of the traditional sources (Chacón, 2005).

Goat's milk is a valid alternative to human's milk, as their nutritional values are largely similar. The taste of goat's milk has little differences compared to that of cows'; they have similar levels of iron, protein, fat, and vitamins C and D. It also has higher contents of potassium, manganese, phosphorous, and vitamins A and B. Doctors and dieticians prescribe this substance as an alternative to bovine milk to people with milk allergies and lactose intolerance. It is also helpful to the elders with intestinal disorders. The international medical corporation certified that goat's milk can revert allergies in children in 50-80% of the cases. Also important is that children who suffer such disorders account for 7% of the world population (Fuenmayor, 2012).

Goat's milk compares to human milk in that it is healthy and nutritious. Many people who suffer from cow's milk allergies may drink goat milk instead, because it contains a different kind of protein (Sánchez, 2011; Bidot *et al.*, 2014).

The protein profile of goat's milk is closer to that of humans, as caprine lactoglobulin is more

easily digested than cow's. Approximately 40% of patients sensitive to the proteins of cow's milk can tolerate goat's milk proteins, because lactalbumin is immunospecific for the two species (Chacón, 2005).

Fat accounts for 3-6% of milk. The quality of caprine milk fat is an important factor, since it defines the capacity of milk for processing. It plays a key role in nutritional and sensorial qualities of by-products (Chávez *et al.*, 2007).

As in other ruminant species, goat's milk fat may be affected by different factors, like breed, individual features, state of lactation, handling, climate, and feed composition. The lipid component is the most important in terms of costs, nutrition, and physical and sensorial features of the product. Within lipids, triglycerides represent about 98%, but goat's milk also has some simple lipids, like diacylglycerol and cholesterol esters, as well as phospholipids and liposoluble compounds, like cholesterol and esters (Park, 2007b).

Richardson(2004) said that the fat of goat's milk is a concentrated source of energy; one unit of it is 2.5-fold more energetic than common carbohydrates. Triglycerides represent almost 95% of total lipids, whereas phospholipids are about 30-40 mg/ml, and cholesterol is 10 mg/100 ml.

The fat composition of goat's milk is the main factor responsible for high cholesterol, because it prevents absorption of excess saturated fats in the organism. Hence, LDL cholesterol and triglycerides are reduced, whereas HDL increases (CAPRAISHISPANA, 2011).

Additionally, goat's milk is characterized by small fat globules (2 µm in goat vs 3-5 µm in cows), which is associated to better digestibility (Alais, 1988; University of Maryland, 1992).

The contents of essential and long chain fatty acids make goat's milk a healthy food for the heart (Capra, 2004). It is also important in the diet of children with atypical forms of eczemas attributed to human's milk with an abnormal profile of fatty acids, especially, linoleic acid (Haenlein, 2002).

#### *Benefits of goat's milk*

Many authors have described the benefits of consuming it (Bello, 1995 a and b; Arbiza, 1996; Haenlein, 2004; Candotti, 2007; Sánchez, 2011). A tenth of the milk consumed in the world comes

from goats, and for some countries, it is the only source of milk (Arbiza, 1987).

The effects of cow's milk allergens on one-year-old children are 3-8% (Maree, 1978; Grezeziak, 1989). The same allergy is observed in 2.5 - 5% of the total world population, with children as the most vulnerable group (Capra, 2004).

In general terms, goat's milk is estimated to supply all the protein an 8 year old child needs, and 6% of proteins needed at 14. In addition to it, each liter of the milk contains 35 g of protein, 54% of the 65g/day required by lactating or pregnant women (Capra, 2004; Candotti, 2007).

Milk is one of the most comprehensive food for humans due to its component traits; proteins contain large amounts of essential aminoacids for nutrition (Paz *et al.*, 2007). Being one of the most complex foods to humans, most countries regard milk production and supply a national priority, and enforce protectionist legislation in this sector (Aréchiga *et al.*, 2008).

Goat's milk demand has increased mainly due to demographic increase and the particular interest in goat's milk by-products in developed countries (yogurt and cheese), because they can be consumed by groups of people who are bovine lactose intolerant. Goat's milk composition is associated to certain nutritional benefits to children, and the development of functional foods and by-products with particular sensorial features required by consumers. This food and its derivatives are another choice to invigorate regional economies (Arbiza, 1996; Haenlein, 2004; Vega and León *et al.*, 2010).

The quality of foods for human consumption largely depends on its potential contribution, both to consumer's sustenance and consumer's health improvement (Es, 1991). As a result, a number of functional foods, nutraceuticals, pharmafoods, modified foods, or substances regarded as nutrients, which can also bring health have been introduced in the market. Also included are designed foods, made to serve a particular purpose or satisfy specific needs of some population groups (Pszczola, 1993; Bello, 1995 a and b), with the common goal of bringing benefits to health. The nutritional composition of goat milk is quite different from other animal-originated foods (Tables 2 and 3), and it is characterized by some nutraceutical traits that make it beneficial to humans. In that sense, it is important to improve the nutri-

tional composition of milk, and sell products with added value. For instance, the nutritional composition of goat milk may be affected by several factors in the diet, including the amount of fiber and the relation between forage and concentrated feeds, which cause major changes, particularly in milk fat (Bedoya *et al.*, 2011; Bidot and Bidot, 2006a).

Goat's milk is a much healthier alternative if consumed whole and from an organic source. The Physiology Department of Granada University has reported that goat's milk has more beneficial properties to humans than cow's milk. The paper was published in the journal *Andalucía Investiga*, and it adds that goat's milk helps prevent ferro-penic anemia (lack of iron) and bone demineralization (osteomalacia). Estimates say that 2% of the milk consumed in the world is produced by goats. Probiotics and other products made from milk are, by far, better than their analogues from cow milk (Solís and Castro, 2007); Flores Córdova *et al.*, 2009).

Beyond the its economic possibilities and use to meet the daily nutritional demands, goat's milk has qualities that make it suitable for children, adults, and lactating mothers (nutraceutical and antiallergic properties). Poorly fed children have seen the benefits of goat's milk as an improved substitute for cow's milk (*Bostaurus*) (Gilbere and Hom, 2002; Capra 2004). However, pediatricians do not recommend it as a complete substitute for human's milk in lactating children under one, due to its high contents of proteins and minerals and low contents of carbohydrates, folic acid and vitamins C, D, E, B6 and B12 (Darnton *et al.*, 1987).

In short, goat's milk is helpful to people with digestive issues (gastric ulcers, gastritis, liver disorders, and intolerance to cow's milk lactose. Regarding allergies, especially those caused by certain bovine dairy proteins, goat's milk has these features,

Similar amounts of sugar and oligosaccharide fractions to human's milk;

A different protein that prevents allergic reactions for consuming cow's milk;

Goat's milk lactose is lower than cow's and human's milk (13% and 41%, respectively). Very low lactose levels make it antiallergic;

Globules or fat drops are smaller and more easily degraded by digestive juices, making it more digestible;

It is more easily tolerated by babies, provided that the mother is unable to breastfeed;

Easier digestion, especially suitable for patients recovering from gastric ailments, ulcers and colitis, thanks to high buffer properties (neutralizes acidity);

The fatty acids contained in goat's milk have the unique capacity to limit cholesterol deposition in body tissues. Cholesterol is lower;

Compared to cow's milk, it has the same amount of proteins, fat, iron, and vitamins C and D. Goat's milk is more abundant in vitamins A and B, and it has lower lactose contents;

It protects against osteoporosis;

It protects against ferro-penic anemia.

## CONCLUSIONS

Due to its traits, biological value as substitute of human's milk, main components and qualities, and the benefits for human health, goat's milk is recommended for children that cannot tolerate other types of milk when they have allergies to particular bovine dairy proteins, as well as for some people with digestive problems, like ulcers, gastritis, liver disorders, cachexia, or are unable to tolerate cow's milk.

## REFERENCES

- ALAIS, C. (1988). *Ciencia de la leche*. México: Continental.
- ARBIZA, A. S. (1996). La leche de cabra. Sus propiedades nutritivas y farmacológicas. *Correo del Maestro*, 3, 1-5.
- ARBIZA, S. I. (1987). *Producción de caprinos*. México: A.G.T.
- ARÉCHIGA, C. F.; AGUILERA, C. A.; RINCÓN, J. I.; MÉNDEZ DE LARA, R. M.; BAÑUELOS S., Meza-Herrera, V. R. (2008). Situación actual y perspectivas de la producción caprina ante el reto de la globalización. *Tropical and Subtropical Agroecosystems*, 9 (1), 1-14.
- BEDOYA, O.; ROSERO, R. y POSADA, SANDRA (2011). *Composición de la leche de cabra y factores nutricionales que afectan el contenido de sus componentes*. Proyecto "Utilización de recursos forrajeros frescos y ensilados, y su impacto sobre la industria láctea caprina", ASOCABRA, Colombia.
- BEDOTTI, F. (2008). *El rol social del ganado caprino*. Retrieved on October 26, 2011, from [http://www.produccionanimal.com.ar/produccion\\_caprina/produccion\\_caprina/11-rol\\_social.pdf](http://www.produccionanimal.com.ar/produccion_caprina/produccion_caprina/11-rol_social.pdf).

- BELLO, J.(1995<sup>a</sup>). Los alimentos funcionales o nutraceuticos. 1. Nueva gama de productos en la industria alimentaria. *Alimentaria*, 26 (1), 25-30.
- BELLO, J. (1995<sup>b</sup>). Los alimentos funcionales o nutraceuticos. 2. Funciones saludables de algunos componentes de los alimentos. *Alimentaria*, 26 (2), 49-58.
- BIDOT, ADELA y MUÑOZ, ROSA (2016). Antecedentes históricos y el origen de las cabras. *Ciencia y Tecnología Ganadera*, 10 (1), 25-30.
- BIDOT, ADELA; SOSA, DANAY y ARTILES, EILÉN (2014). Importancia de la leche de cabra en la alimentación humana. Nota divulgativa. *Ciencia y Tecnología Ganadera*, 8(3), 175-178.
- BIDOT, ADELA y BIDOT, G. (2006<sup>a</sup>). La producción de leche caprina y sus formas de comercialización. Recopilación. *Revista Agroenfoque*, 3 (1), 21-25.
- BIDOT, ADELA (2006<sup>b</sup>). *La cabra como productora de leche*. Memorias CD III Congreso Internacional sobre Mejoramiento Animal, La Habana, Cuba.
- BORRAS, A. (1968). *Cómo comer y beber leche*. La Habana, Cuba: Comité Nacional Lechero.
- BOZA, J. (1981). *Mejora de la cabra granadina*. España: Caja Provincial de Ahorros de Granada.
- BRENNEMAN, J. C. (1978). *Basics of Food Allergy*. Springfield, Illinois, EE.UU.: Ch. C Thomas Publishing.
- CANDOTTI, J. J. (2007). *Los beneficios de la leche caprina en la infancia*. Retrieved on February 10, 2016, from [www.todoagro.com.ar](http://www.todoagro.com.ar).
- CAPDEVILA, F. y MARTÍ-HENNEBERG, C. (1996). Trascendencia nutricional del consumo de lácteos en la dieta mediterránea actual en España. *Alim. Nutri. Salud*, 3, 9-17.
- CAPRA (2004). *La composición de la leche de cabra y su papel en la alimentación humana* (en línea). Retrieved on April 1, 2016, from <http://www.iespana.es/CAPRA/HOMBRE/HOMBRE.HTM>.
- CAPRAISPANA (2011). *La composición de la leche de cabra y su papel en la alimentación humana*. Retrieved on April 1, 2016, from <http://www.capraispana.com/destacados/hombre/hombre.htm>.
- CHACÓN, A. (2005). *Aspectos nutricionales de la leche de cabra (Caprahircus) y sus variaciones en el proceso agroindustrial*. Retrieved on April 8, 2016, from [http://www.mag.go.cr/rev\\_meso/v16n02\\_239.pdf](http://www.mag.go.cr/rev_meso/v16n02_239.pdf).
- CHÁVEZ, M. S.; MARGALEF, M. I.; MARTÍNEZ, M. (2007). *Cuantificación de lipólisis en leche caprina (Saanen) cruda y térmicamente tratada*. Retrieved on April 8, 2016, from [http://www.produccionanimal.om.ar/produccion\\_caprina/leche\\_caprina/38-Cuantificlipolisis-leche.pdf](http://www.produccionanimal.om.ar/produccion_caprina/leche_caprina/38-Cuantificlipolisis-leche.pdf).
- CHILLIARD, Y.; FERLAY, A.; ROUEL, J.; LAMBERET, G. (2003). A Review of Nutritional and Physiological Factors Affecting Goat Milk Lipid Synthesis and Lipolysis. *Journal of Dairy Science*, 86, 1751-1770.
- COFRÉB. P. (2001). Producción de cabras lecheras. Chillán. *Boletín INIA*, 66, 132-134.
- CRUZ, ANITA; MOSQUERA, J. N. y CLAVIJO, M. (2012). *Caracterización de sistemas de producción de leche caprina en el sur del Uruguay*. Tesis en opción del título de Ingeniero Agrónomo. Universidad de la República, Facultad de Agronomía, Montevideo, Uruguay.
- DARNTON, I.; COVENEY, J.; DAVEY, G. R. (1987). Goat Milk, Nutritional and Public Health Aspects: a Review. *Food Technology in Australia*, 39 (12), 572-688.
- DORIA, S. (1997). *Caprinocultura, cría racional de caprinos*. San Pablo, Brasil: Livraria Nobel.
- EDWARD, F. (2012). *Los beneficios de la leche de cabra; ¿una alternativa a la leche de vaca?* EE.UU.: Global Healing Center.
- VAN, E. (1991). *Animal Nutrition and Human Health. Lecture of Prize Roche Research for Animal Nutrition*. Retrieved on March 28, 2013, from <http://om.iamm.fr/om/pdf/a67/06600037.pdf>.
- FAO (2010). *Greenhouse Gas Emissions from the Dairy Sector*. Retrieved on March 28, 2013, from <http://www.fao.org/docrep/012/k7930e/k7930e00.pdf>.
- FAO (1987). *Tecnología de la producción caprina*. Santiago de Chile.
- FAO (1982). *Anuario FAO de Producción*, 36, 217-219.
- FLORES-CÓRDOVA, M. A.; PÉREZ-LEAL, R., BASURTO-SOTELO, M. y JURADO-GUERRA, M. R. (2009). La leche de cabra y su importancia en la nutrición. *TECNOCENCIA Chihuahua*, 3 (2), 107-113.
- FUENMAYOR, R. (2012). *Comunicación personal*, 12 de enero de 2012.
- GILBERE, G. y HOM, D.A. (2002). *The Magic of Goat Milk*. Retrieved on April 1, 2016, from [http://fredompressonline/FPO\\_feacturedArticles\\_carpa.htm](http://fredompressonline/FPO_feacturedArticles_carpa.htm).
- GREZESIAK, T. (1989). Prescription of Goat Milk in Pediatrics-Revolutionary? *Le Concours Medical*, 111, 3059-3064.
- HAENLEIN, W. (2004). Goat Milk in Human Nutrition. *Small Ruminant Research*, 51, 155-163.
- HAENLEIN, W. (2002). *Milk and Meat Products*. Retrieved on August 19, 2015, from [http://goatconnection.com/articles/publish/article\\_73.shtml](http://goatconnection.com/articles/publish/article_73.shtml).
- JANDAL, J. M. (1996). Comparative Aspects of Goat and Sheep Milk. *Small Rumin Res.*, 22, 177-185

- KNIGHTS, M.; GARCÍA, G. W. (1997). The Status and Characteristics of the Goat (*Capra hircus*) and its Potential Role as a Significant Milk Producer in the Tropics, A Review. *Small Ruminant Research*, 26 (3), 203-215.
- LÓPEZ, J. C.; FUENTES, V. H.; FIGUEROA, J. J.; SÁNCHEZ, R. A.; SERNA, A.; RUIZ, J. I.; *et al.* (2011). *Técnicas para la transformación de leche de cabra en zonas marginales*. México: Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación.
- MAREE, H. P. (1978). Goat Milk Anits Use as Hypo-Allergenic Infant Food. *Dairy Goat Journal*, 43, 363-365.
- MENESES, R. R. (2007). Manejo de ordeño para mejorar la calidad de la leche de cabra. Chile: INIA.
- ORTEGA, G.; RAZ, I.; MAGAÑA, H.; ORTIZ, J.; SIERRA, S.; CENTURIÓN, F. *et al.* (2011). Interacción genotipo x ambiente en cabras lecheras. *Bioagrociencias*, 4 (2), 32-40.
- OTTOGALLI, G. y TESTOLIN, G. (1991). Dairy Products. En G. A. Spillered, A. V. y Van Nostrand (eds) *The Mediterranean Diet in Health And Disease* (pp. 135-139). Nueva York: Botanical Garden Press.
- PARK, Y. W.; JUÁREZ, M.; RAMOS, M. y HAENLEIN, G. (2007). Physico-Chemical Characteristics of Goat and Sheep Milk. *Small Ruminant Research*, 68, 88-113.
- PARK, Y. M. (2007a). Rheological Characteristic of Goat and Sheep Milk. *Small Ruminant Research*, 68, 73-87.
- PARK, Y. W. (2007b). Phisico-chemical Characteristics of Goat and Sheep Milk, *Small Ruminant Research*, 68, 88-113.
- PARK, Y. W. (2006). Goat Milk Chemistry and Nutrition. En Y. W. Park y F. W. Haenlein (Eds.), *Handbook of Milk of Non-bovine Mammals* (pp. 34-58). Blackwell Publishing Professional, Oxford, UK/Ames, Iowa.
- PAZ, R. G.; TOGO, J. A. y LÓPEZ, C. (2007). *Evaluación de parámetros de producción de leche en caprinos* (Santiago del Estero, Argentina). *Revista Científica FCV-LUZ*, 17 (2), 161-165.
- PSZCZOLA, D. E. (1993). Alimentos de diseño: un concepto que evoluciona. *Alimentaria*, 22 (1), 91-93.
- RICHARDSON, C. W. (2004). Let's Learn About Dairy Goats and Goat's Milk. Oklahoma Cooperative Extension Service Oklahoma State University. *Dairy Goat Association Bulletin*, (424), 1-4. Retrieved on March 21, 2013, from <http://oklahoma4h.okstate.edu/litol/file/backup/4h424.pdf>.
- RODRÍGUEZ, C. A. y VALENCIA, Ch. E. (2006). La leche de caprina, otras propiedades y atributos. *Agricultura*, 2(1), 4-6.
- SAINI, A. L.; GILI, R. S. (1991). Goat Milk: an Attractive Alternative. *Indian Dairyman*, 42, 562-564
- SALVADOR, A.; MARTÍNEZ, G.; ALVARADO, C. y HAHN, M. (2006). Composición de la leche de cabras mestizas Canarias en condiciones tropicales. *Zootecnia Tropical*, 24 (3), 12-16.
- SÁNCHEZ, M. (2011). *La leche de cabra tiene los mismos nutrientes que la materna sin ser alérgica*. España: Universidad de Granada.
- SINGH, V. B. y SINGH, S. N. (1985). Amino-acid Composition of Casein of Four Indian Goat-Breeds During Lactation. *Asian Journal of Dairy Research*, 3 (4), 187-192.
- SOLÍS, R. J. y CASTRO, R. A. (2007). La leche de cabra en la nutrición y en la terapéutica. *Revista de la Universidad de Chapingo*, 4(1), 22-47.
- UNIVERSITY OF MARYLAND (1992). *Nacional Goat Handbook*. Retrieved on November 16, 2014, from <http://www.inform.umd.edu/EdRes/topic/AgrEnv/ndd/goat>.
- VACAS, C. (2003). *Evolución del sector caprino en la Región de Murcia (1986-2000) y su caracterización productiva al final del milenio*. Retrieved on March 24, 2013, from <http://www.tesisenred.net/handle/10803/11020>.
- VARGAS, P.; PINEDA, M. I.; CHACÓN, A. (2007). Lácteos bovinos y percepción de la leche caprina entre estudiantes de la Universidad de Costa Rica. *Agronomía Mesoamericana*, 18 (1), 27-36.
- VEGA, Y.; LEÓN, S.; GUTIÉRREZ, R.; DÍAZ, G.; GONZÁLEZ, M.; RAMÍREZ, A., *et al.* (2010). *Leche de cabra: producción, composición y aptitud industrial*. Retrieved on March 1, 2015, from <http://www.alfaeditores.com/carnilac/TECNOLOGIA%20Leche%20de%20cabra.html>.

Received: 1-12-2017

Accepted: 1-20-2017

**Table 1. Composition of goat's milk**

Composition of goat milk (%)	
Total solids	11.70-15.21
Protein (Nx6,38)	2.90-4.60
Fat	3.00-6.63
Lactose	3.80-5.12
Ashes	0.69-0.89
pH	6.41-6.70

Source: Boza *et al.*, 1992, cited by Cruz *et al.*, 2012

**Table 2. Comparison of milk composition in different species (%)**

Component	Goat	Sheep	Cow	Woman
Water (%)	86.20	80.90	87.50	88.35
Fat (%)	3.80	7.62	3.67	3.67-4.70
Nonfatty solids (%)	8.68	10.33	9.02	8.90
Lactose (%)	4.08	3.7	4.78	6.92
Protein (%)	2.90	6.21	3.23	1.10
Casein (%)	2.47	5.16	2.63	0.40
Serum proteins (%)	0.43	0.81	0.60	0.70
Ashes (%)	0.79	0.90	0.73	0.31
Vitamin A (IU)	185	146	126	190
Vitamin D (IU)	2.3	0.18	2.0	1.4
Thiamin (mg)	0.068	0.08	0.045	0.017
Riboflavin (mg)	0.21	0.376	0.16	0.02
Niacin (mg)	0.27	0.41	0.08	0.17
Pantotenic acid (mg)	0.31	0.408	0.32	0.20
Vitamin B6	0.046	0.08	0.042	0.011
Folic acid	1.0	5.0	5.0	5.5
Biotin (µg)	1.5	0.93	2.0	0.4
Vitamin B12	0.065	0,712	0.357	0.03
Vitamin C	1.12	4.16	0.94	5.00
Energy (cal/100 ml)	70.00	Nd	69.00	68.00

Source: Jandalet *et al.*(1996)



**Table 3. Comparison of three types of milk, composición in 100 mL**

	Human	Cow	Goat
Protein (g)	1.2	3.3	3.3
Casein (g)	0.4	2.8	2.5
Lactoalbumin (g)	0.3	0.4	0.4
Fat (g)	3.8	3.7	4.1
Lactose (g)	7.0	4.8	3.8
Calorific value (Kcal)	71	69	76

Source: CAPRAHISPANA (2011)

**Table 4. Mineral contents in cow's and goat's milk(100 g)**

Component	Goat	Cow
Ca (mg)	134	122
P (mg)	121	119
Mg (mg)	16	12
K (mg)	181	152
Na (mg)	41	58
Cl (mg)	150	100
S (mg)	28	32
Fe (mg)	0.07	0.08
Cu (mg)	0.05	0.06
Mn (mg)	0.032	0.02
Zn (mg)	0.56	0.53
I (mg)	0.022	0-021

Source: Park (2006)